



IJBEM 1999, 1(1), 47-53 • www.tut.fi/ijbem/

Clinical Application of MRI Image Processing in Neurology

Frey H^a, Lahtinen A^b, Heinonen T^b, Dastidar P^c

^a Department of Neurology, Tampere University Hospital, Tampere, Finland

^b RagnarGranit Institute, Tampere University of Technology, Tampere, Finland

^c Department of Radiology, Tampere University Hospital, Tampere, Finland

Abstract

The development of new computer based medical imaging methods have made the extremely rapid progress in both basic and clinical neurosciences possible. Among the new techniques MRI has made quantum leaps during the eighties and nineties and has the greatest promise for future development especially when combined by fusion with other techniques to make the multimodal imaging and mapping of both the structure and function of the brain possible. In the following overview only a few of these developments can be partially covered. In order to understand the real influence of these technical advances several recent textbooks and thousands of individual articles plus abstracts should be summarized. Therefore also the prediction of the future progress of the most promising avenues to find new and relevant information is made impossible. A clinician can only present some ideas of the existing needs.

1. Remarks on Advances of Radiological Techniques in Neurology

Radiological techniques started to contribute relatively late in clinical neurology after the original discovery of x – rays in 1895 by Roentgen. The application of conventional skull x-ray techniques improved the diagnosis and treatment of skull traumas. But first the development of pneumoencephalography for the study of the soft tissues and ventricles in the brain and myelography after injection of air or inert gas into the CSF space started a new stage of development. These techniques were slowly substituted after late fifties by the introduction of contrast media (aorticervical and later selective angiographies and water insoluble and later water soluble opaque contrastmedia for myelography).

A new and revolutionary stage of the imaging techniques in neuroradiology and neurology started, when Hounsfield and Cormack developed CT (computerized tomography) and received a Nobel price in 1972. CT uses ionizing radiation, but it has no other major risks and is relatively comfortable for the patient. Until the development of spiral- techniques and 3D modelling the CT slices have been limited primarily to transaxial views. In CT images are generated by passing an x–ray beam through the skull or other object (e.g. spine and vertebral column) and measuring its degree of attenuation. The ability to attenuate x – rays of different tissues (e.g. brain, bone and CSF differs and can be measured numerically as a tissue density number for each voxel (volume element). These numbers can be converted to gray scale values and presented visually as pixels. A large number of reconstruction and

In dementia of Alzheimer type the neurodegenerative changes particularly in hippocampus (volumetry) and related areas in temporal and parietal cortex, where also neuropathological postmortem changes are usually most profound help to make the right diagnosis. More recently such dementia entities as frontal lobe dementia and Lewy body dementia have been added among the differential diagnostics based partially on MRI. In Huntington's disease patterns of hypometabolic activity are found in the basal ganglia of both the patients and their relatives in risk. FMRI can in near future be used in the diagnosis of Parkinson's disease and other extrapyramidal disorders. Recent findings of structural changes seen in MRI in schizophrenia are giving new light to the possible underlying pathology and probable early developmental lesions in this disease.

3. Recent Advances in the Development and Application of Neuroimaging Methods in Neurology

The increasing use of different computer based imaging methods, multimodal imaging and 3D modelling and fusion of different imaging methods with each others have made new developments and application for use also in clinical applications possible. For these new developments are responsible especially the research initiatives, which have emerged after the publication of the Mapping the Brain and its functions report in U.S.A. (Poscura P and Martin J, 1992). For overviews in the recent developments see e.g. Toga AW and Mazziotta JG 1996, Frackowiak RSJ et al. 1997.

3.1. Digital neuroatlases and their application

There exist several different ways and tools for building human (or animal) brain atlases. Atlas is a special form of map that incorporates stereotactic elements. Computerized stereotactic atlases are becoming more important for the localization and segmentation of human brain anatomy and physiology. Some of the tools available to analyse brain structures and functions at macroscopic level are listed in Table I.

Table I. Tools for macroscopic study of the brain

CT
MRI
MR spectroscopy (combined with topography)
SPECT
PET
EEG
MEG
Evoked potentials
Postmortem cryosectioning combined with staining
Multimodal imaging

The most straightward approach is based on the use of only one fixed parameter for spatial normalization. E.g. shape and adjust the size (MRI image or section). Here in comparisons between different approaches the so-called densitometric approach seems to be the most reliable method sofar. Instead of this Talairach and Tournoux used anterior and posterior brain commissures as reference points.

Segmentation is a third alternative e.g. to separate white matter from gray matter or ventricles (CSF space). Brain image segmentation is especially well suited for this type of an approach.

Visualization is helpful especially in multimodal imaging atlases (fusions), where different colours can be painted over a grey scale.

The probabilistic approach means, that a large population of samples is treated statistically and medians of these digital samples are represented as standards. E.g. at the Montreal Neurological Institute a MRI

brain atlas was created by combining MRI images taken from 305 normal 28 year old individuals. After this a median stereotactic digital neuroatlas was developed. (Evans AC et al. 1992).

One has to keep in mind, that a brain atlas is essentially also a database. By combining the other existing information with different types of databases (books, articles etc.) it is possible to increase enormously the efficiency of a digital neuroatlas.

We in Tampere have developed a different type of an approach. The original digital neuroatlas is transferred on the corresponding sections of the MRI image of the patient. The shape and size of the neuroatlas can be modified by using different mathematical algorithms (rubber sheet and wrap algorithms). Also the possible errors in the patient image (rotations, translations and stretching) can be corrected (Lahtinen A et al 1998). Some of the neuroatlases, which can be found in Internet are listed in Table II.

Table II. Neuroatlases in Internet

Talairach – Tournoux Brain Atlas
Talairach – Tournoux Referential Brain Atlas
Brodman Gyri Brain Atlas
SchaltenBrand – Wahren Brain Atlas (axial-, coronal-, sagittal slices.
Ono Kubik Sulcal Atlas (indexed)
The whole brain atlas
Atlas at Washington
Ito´s Brain Atlas
PET Brain Atlas
Atlas of Brain perfusion SPECT
The Human Brain: Dissections of the Real Brain




Figure 2. An example of the Tampere Brain Atlas to illustrate a brain infarct in the right *a.cerebri media* lesioning inferior frontal gyrus, both precentral and postcentral gyri and inferior parietal gyrus.

A second example of the use of neuroatlas, which shows an brain haemorrhage in a 3D axial serial section image




Figure 3. 3D image of a large haemorrhage. This type of image can be used e.g. in planning of neurosurgery.

A third example is the localization of multiple sclerosis plaques in the white matter of brain after segmentation (Fig 4). This type of an approach is currently used to follow the possible effect of beta-interferon treatment of MS patients (Dastidar P. et al. 1999).




Figure 4. Axial T1 and T2 weighted MRI sections of MS plaques and brain atrophy.

However there are still problems transforming brains to a common atlas. Because of the individual variance and changes due to ageing the accuracy of brain atlases is restricted to limited regions of the Brain. Especially cortex is problematic in this sense (Benayoun et al. 1994). Subcortical areas are more homologous and the points here are more easily identifiable (Kazanovskaya et al. 1991).

3.2 Application of fMRI in Clinical Neurology

fMRI techniques are noninvasive, multiple, longitudinal. When compared with other functional brain imaging techniques (EEG, PET, magnetic source imaging and near infrared spectroscopic imaging, fMRI has good temporal resolution and excellent spatial resolution. Therefore it has become a new and very powerful research tool (Karni, A. et al. 1995) and has started to show considerable clinical benefit while studying the underlying pathology in different brain diseases. In future it may also be used to follow up the recovery of the brain lesions and be used to help in restorative neurology for higher cognitive functions.

In most fMRI studies, image sets are acquired while the patient is alternatively in an active and control state. While changes in cerebral blood flow and its oxygenation degree are measured. In the near future these types of techniques can also be applied to study the possible regional pathological changes related to CBF. In this connection fMRI has to be compared with diffusion imaging EPI techniques and SPECT or PET.

3.3. Clinical Applications of Volumetric image Analysis in Neurology

Despite of the still existing difficulties in practical application of volumetry these techniques have already been set widely to study such degenerative diseases of the brain as Alzheimer-type dementia (Erkinjuntti et al. 1993, Devernoy 1998). Brain tumors (Velthuisen et al.1995), brain infarcts (Heinonen et al.1998) and intracerebral hematomas (Dastidar et al.1997) aswellas MS plaques (Dastidar et al.1999).

In the near future 3D visualization will become a valuable tool for this type of applications in clinical neurology and planning of stereotactic neurosurgery, radiotherapy and follow up of the effect of treatment. (Heinonen et al.1998).

3. Some Future Prospects and needs

Especially in clinical neurosurgery the use of stereotactic methods, when combined with 3D modelling would benefit the planning and surgical treatment immensely. The same is due to radiotherapy of brain tumors. The future applications of fMRI and multimodal imaging techniques will open up totally new ways to understand the cognitive and psychiatric disorders. The combination of these techniques with already existing international databases and their use will also improve the diagnostics and treatment of brain and other nervous system disorders much. At the same time the need to educate new clinical neuroscientists and clinicians who can understand and use these techniques will become an absolute necessity. In fact it is necessary to train a new generation of specialists to fulfil all these demands.

References

1. Osborn A.G.: Diagnostic neuroradiology. Mosby Press.Mosby – Year Book Inc.1994.
2. Gonzalez et al., Head & spine imaging. Wiley Medical Press.1985.
3. Taveras, J.M., Neuroradiology, Third ed. Williams and Wilkins Press.1996.
4. Bradley W.G.Jr. and Bydder G.M., Advanced MR Imaging techniques. Martin Dunitz LTD, London. 1997.
5. Talairach J. and Tournoux P. Co – planar stereotaxic atlas of the human brain. 1988. Thieme, New York.
6. Evans AC, Collins DI and Milner B., An MRI – based stereotactic brain atlas from 300 young normal subjects. Proc. 22nd Ann.symposium of the society of neuroscientists, Anaheim, 1992.
7. Poscura J. and Martin J.Mapping the human brain and its functions. Institute of Medicine. 1992.
8. Toga AW and Mazziotta JC. Brain mapping, the methods. Academic Press.1996.
9. Frackowiak RSJ, Friston, KJ.,Frith CD, Dolan RJ and Mazziotta JC.: Human brain function. Academic Press 1997.
10. Lahtinen A.,Kulkas,T.,Heinonen T. and Frey H.: The application of a new digital neuroatlas for post-stroke dementias. Norage 1998 Proc. 86,1998.
11. Dastidar P., Heinonen T. Vahvelainen T.Elovaara I and Eskola H.: Computerized volumetric analysis of lesions in multiple sclerosis using a new semiautomatic segmentation software. Med. Biol. Engineering & computing. 37: 104 – 107. 1999.
12. Benayoun S., Ayache N. and Cohen I.: An adaptive model for 2D and 3D dense non-rigid motion computation. INRIA Inter.Rep. No 2297, 1994.
13. Kazarnovskaya MI., Borodkin SM., Shablov VA., Krivosheina VY and Golanov AV.: 3 – D computer model of subcortical structures of human brain. Comput.Biol.Med. 21(6),451-457, 1991.
14. Karni A., Meyer P., Jezard M, Adams M, Turner R. and Ungerleider LG.: Functional MRI evidence for adult motor cortex plasticity during motor skill learning. Nature 377,155-158 (1995)
15. Erkinjuntti T.,Lee D.,Gao F.,Steenhuis R.,Eliasziw M.,Fry R.,Merskey H and Hachinski V.: Temporal lobe atrophy on magnetic resonance imaging in the diagnosis of early Alzheimer's disease. Arch.Neurol.50, 305-310.1993.
16. Duvernoy H.: The human hippocampus.2.ed. Springer Verlag Berlin Heidelberg.1998.
17. Velthuisen R.,Clarke L.,Phuphanics S.,Hall O.,Bensaid A.,Arrington J.,Greenberg HM. And Silbiger M.: Unsupervised measurement of brain tumor volume on MR images.J.Magnetic Resonance Imaging.5.594-605.1995.
18. Heinonen T, Dastidar P.Eskola H., Frey H., Ryymin P and Laasonen E.: Applicability of semi-automatic segmentation for volumetric analysis of brain lesions.J.Med.Eng.Technol.22.173-178.1998.
19. Dastidar P, Heinonen T., Virta T., Ryymin P. and Kuurne T.: Evaluation of operative treatment of intracerebral hematomas by semiautomatic segmentation method. Proc. 9th int.conference.Biomed.Engineering.Singapore.733.1997.
20. Heinonen T., Visala K., Blomqvist M., Eskola H. and Frey H.: 3Dvisualization library for multimodal images. Comp.Med.Imaging and Graphics.22.267-273.1998.

Clinical applications of functional magnetic resonance imaging. *Neuroimaging Clin N Am* 2001;11:221-36, viii. [OpenUrl PubMed](#).
Stippich C, Freitag P, Kassubek J, et al. Motor, somatosensory and auditory cortex localization by fMRI and MEG. *Neuroreport* 1998;9:1953-7.
Potentially adaptive functional changes of cognitive processing in patients with multiple sclerosis and their acute modulation by rivastigmine. *Brain*, (in press).
Ward NS, Brown MM, Thompson AJ, et al. Neural correlates of outcome after stroke: a cross-sectional fMRI study. *Brain* 2003;126:1430-48. [OpenUrl Abstract/FREE Full Text](#).
High-resolution magnetic resonance imaging (HR-MRI) is a noninvasive method for the diagnosis of VBA-DA. VBA-DAs were classified according to the feature of HR-MRI in combination with digital subtraction angiography (DSA), and the clinical outcomes of different types of VBA-DAs were analyzed.
2.2 High-resolution magnetic resonance imaging. All patients underwent HR-MRI scanning with a 3.0T GE device (HDx 3.0 T, GE Healthcare, Chicago, IL). MRI scans were performed in 8 channel head and neck coils with a matrix size of 256 × 256 pixels and a pixel size of 0.28 mm × 0.28 mm. The lesion region was identified by 3D-TOF-MRA. Two experienced neuroradiologists independently reviewed the images on post-processing workstation in a double-blind way.